**Harbor-UCLA Interfacility Report**
Davita Fax: 310-533-1709

Date:

Patient Name: D.O.B:

Type of Infection: ☐ Peritonitis ☐ Exit site ☐ Tunnel

1- Duration of therapy: Therapy in hospital:

2- Nystatin 5 ml swish and swallow 1 week beyond the course of antibiotic therapy
3- Choice of Intraperitoneal Antibiotic at the time of discharge for continuous PD:
(Please indicate the total dosage in front of each option)
((LD = loading dose in mg/L; MD =maintenance dose in mg/L; all exchanges.)

☐ Gentamicin, netilmicin, or tobramycin (LD 8, MD 4):

☐ Cefazolin, cephalothin, or cephradine (LD 500, MD 125):

☐ Cefepime (LD 500, MD 125):

☐ Ceftazidime (LD 500, MD 125):

☐ Vancomycin (LD 1000, MD 25):

☐ Amoxicillin (LD 250–500, MD 50):

☐ Ampicillin, oxacillin, or nafcillin (MD 125):

☐ Imipenem/cilastin (LD 250, MD 50):

☐ Ciprofloxacin (LD 50, MD 25):

☐ Aztreonam (LD 1000, MD 250):

☐ Daptomycin (LD 100, MD 20):

☐ Linezolid Oral 200–300 mg q.d.

☐ Ampicillin/sulbactam (LD 1000, MD 100):

☐ Trimethoprim/sulfamethoxazole Oral 960 mg b.i.d

☐ Other:

4- Result of peritoneal Culture: ☐ No growth ☐ Organism:

5- Sensitivity: ☐ Date: ☐ at Harbor ☐ at Davita

6- Follow up Appointment: The day after discharge

Discharge Fellow: